



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 233/25, 237/36, 237/40, A61K 47/18	A2	(11) International Publication Number: WO 00/07979 (43) International Publication Date: 17 February 2000 (17.02.00)												
(21) International Application Number: PCT/US99/17974 (22) International Filing Date: 6 August 1999 (06.08.99) (30) Priority Data: <table border="0"> <tr> <td>60/095,778</td> <td>7 August 1998 (07.08.98)</td> <td>US</td> </tr> <tr> <td>60/098,500</td> <td>31 August 1998 (31.08.98)</td> <td>US</td> </tr> <tr> <td>60/108,366</td> <td>13 November 1998 (13.11.98)</td> <td>US</td> </tr> <tr> <td>60/119,207</td> <td>5 February 1999 (05.02.99)</td> <td>US</td> </tr> </table> (71) Applicant (for all designated States except US): EMISPHERE TECHNOLOGIES, INC. [US/US]; 765 Old Saw Mill River Road, Tarrytown, NY 10591 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GSCHNEIDNER, David [US/US]; 44 Cerretta Street #6, Stamford, CT 06907 (US). LEONE-BAY, Andrea [US/US]; 20 Woodland Way, Ridgefield, CT 06877 (US). WANG, Eric [US/US]; 8205 Featherhill Road #302, Perryhall, MD 21128 (US). ERRIGO, Lynn [US/US]; 85 Birch Street, Port Chester, NY 10573 (US). KRAFT, Kelly [US/US]; 38 1/2 Wolden Road, Ossining, NY 10562 (US). MOYE-SHERMAN, Destardi [US/US]; 102 Chateau Rive, Peekskill, NY 10566 (US). HO, Koc-Kan [MC/US]; 70 Barker Street, Mount Kisco, NY 10549 (US). PRESS, Jeffrey, Bruce [US/US]; Bear Berry Lane, Brew-		60/095,778	7 August 1998 (07.08.98)	US	60/098,500	31 August 1998 (31.08.98)	US	60/108,366	13 November 1998 (13.11.98)	US	60/119,207	5 February 1999 (05.02.99)	US	ster, NY 10509 (US). WANG, Nai, Fang [CN/US]; Apartment 2C, 41-42 42nd Street, Long Island City, NY 11104 (US). (74) Agents: LESSLER, Jay, P. et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022-7513 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
60/095,778	7 August 1998 (07.08.98)	US												
60/098,500	31 August 1998 (31.08.98)	US												
60/108,366	13 November 1998 (13.11.98)	US												
60/119,207	5 February 1999 (05.02.99)	US												
(54) Title: COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS														
(57) Abstract <p>Compounds and compositions for the delivery of active agents are provided. Methods of administration and preparation are provided as well.</p>														

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS

FIELD OF THE INVENTION

The present invention relates to compounds for delivering active agents, such as biologically or chemically active agents, to a target. These compounds are well suited for forming non-covalent mixtures with active agents for oral, intracolonic, or other routes of administration to animals. Methods for the preparation and administration of such compositions are also disclosed.

BACKGROUND OF THE INVENTION

Conventional means for delivering active agents are often severely limited by biological, chemical, and physical barriers.

Typically, these barriers are imposed by the environment through which delivery occurs, the environment of the target for delivery, and/or the target itself. Biologically and chemically active agents are particularly vulnerable to such barriers.

In the delivery to animals of biologically active and chemically active pharmacological and therapeutic agents, barriers are imposed by the body. Examples of physical barriers are the skin, lipid bi-layers and various organ membranes that

are relatively impermeable to certain active agents but must be traversed before reaching a target, such as the circulatory system. Chemical barriers include, but are not limited to, pH variations in the gastrointestinal (GI) tract and degrading enzymes.

These barriers are of particular significance in the design of oral delivery systems. Oral delivery of many biologically or chemically active agents would be the route of choice for administration to animals if not for biological, chemical, and physical barriers. Among the numerous agents which are not typically amenable to oral administration are biologically or chemically active peptides, such as calcitonin and insulin; polysaccharides, and in particular mucopolysaccharides including, but not limited to, heparin; heparinoids; antibiotics; and other organic substances. These agents may be rapidly rendered ineffective or destroyed in the gastro-intestinal tract by acid hydrolysis, enzymes, and the like. In addition, the size and structure of macromolecular drugs may prohibit absorption.

Earlier methods for orally administering vulnerable pharmacological agents have relied on the co-administration of adjuvants (e.g., resorcinols and non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to increase artificially the permeability of the intestinal walls, as well as the co-administration of enzymatic inhibitors (e.g., pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) and trasylol) to inhibit enzymatic degradation. Liposomes have also been described as drug delivery systems for insulin and heparin. See, for example, US 4,239,754; Patel et al. (1976), *FEBS Letters*, Vol. 62, pg. 60; and Hashimoto et al. (1979), *Endocrinology Japan*, Vol. 26, pg. 337. However, broad spectrum use of such drug delivery systems is precluded because: (1) the systems require toxic amounts of adjuvants or inhibitors; (2) suitable low molecular weight cargos, i.e. active agents, are not


available; (3) the systems exhibit poor stability and inadequate shelf life; (4) the systems are difficult to manufacture; (5) the systems fail to protect the active agent (cargo); (6) the systems adversely alter the active agent; or (7) the systems fail to allow or promote absorption of the active agent.

More recently, microspheres of artificial polymers of mixed amino acids (proteinoids) have been used to deliver pharmaceuticals. For example, US Patent 4,925,673 describes drug-containing proteinoid microsphere compounds as well as methods for their preparation and use. These proteinoid microspheres are useful for the delivery of a number of active agents. In addition, certain modified amino acids have been used to deliver pharmaceuticals. See, e.g., US 5,629,020; US 5,643,957; US 5,650,386; and US 5,776,888.


However, there is still a need for simple, inexpensive delivery systems which are easily prepared and which can deliver a broad range of active agents by various routes.

SUMMARY OF THE INVENTION


Compounds and compositions which are useful in the delivery of active agents are provided. The compounds comprise the following compounds or salts thereof.



<u>Cpd. #</u>	<u>n</u>	<u>m</u>	<u>X</u>
1	3	CH ₂ O	2-OH
2	3	CH ₂ O	4-OH
3	3	0	2-NH ₂ , 5-F
4	2	0	2-NH ₂ , 5-F
5	3	0	2-NH ₂ , 5-Cl
6	2	0	2-NH ₂ , 3,5-Cl
7	2	0	2-NHMe
8	3	1	4-OH
9	3	1	3-OH
10	3	0	2-NHMe
11	2	0	2-OH, 3-F; 5-Cl
12	2	0	2-OH, 3-Cl; 5-F
13	2	0	2-OH, 3,5-Me
14	3	0	2-OH, 3,5-Me
15	2	0	2-OH, 3-Br, 5-Cl
16	3	2	2-OH
17	2	2	2-OH
18	2	0	2-OH, 3,5-F
19	3	0	2-OH, 3,5-F
20	2	0	2-OH, 5-F
21	3	0	2-OH, 5-F
22	2	0	2-NHAc
23	3	0	2-SO ₃ Na
24	3	0	2-OH, 3-Me, 5-F
25	3	0	2-OH, 3-Me, 5-Cl
104	3	0	2-OH, 4-Ome
133	2	0	2-OH, 3-Me, 5-Cl




<u>Cpd#</u>	<u>n</u>	<u>X</u>
26	7	2-OH, 5-Me
27	6	2-OH
28	8	2-OH, 3,5-Cl
29	7	2-OH, 3,5-Cl
30	8	2-OH, 4-Me
31	7	2-CH ₂ OH
32	4	2-OH, 4-Me
33	7	2-OH, 4-Me
34	7	2-OH, 5-F
35	8	2-OH, 5-F
119	3	2-OH, 5-Cl
120	5	2-OH, 5-Cl
121	6	2-OH, 5-Cl
122	7	2-OH, 5-Cl
123	8	2-OH, 5-Cl
124	1	2-OH, 5-Cl
125	2	2-OH, 5-Cl
126	4	2-OH, 5-Cl
127	9	2-OH, 5-Cl
128	10	2-OH, 5-Cl
129	11	2-OH, 5-Cl
130	12	2-OH, 5-Cl
131	7	2-OH, 3,4-F
132	7	2-OH, 4-F




Cpd #	n	m	X
36	7	0	2-OH, 3-NH ₂ , 5-NO ₂
37	5	0	2-OH, 4-Cl
38	7	CH ₂ O	4-OH
39	7	0	2-NH ₂ , 5-F
40	7	0	2-NH ₂ , 5-Cl
41	7	0	2-OH, 3,5-F
42	7	0	2-OH, 3,4-F
43	7	0	2-NHMe
44	7	0	2-OH, 4-F
45	7	0	2-OH, 3-F, 5-Cl
46	7	1	4-OH
47	7	0	2-OH, 3-Cl, 5-F
48	7	0	2-OH, 3-Br, 5-Cl
49	7	0	2-OH, 3,5-Me
50	7	0	2-OMe, 6-Cl
51	7	0	2-OH, 6-Cl
52	7	1	3-OH
53	7	2	2-OH
54	7	0	2-OH, 5-F
55	7	0	2-OH, 3-Me, 5-Cl
56	7	0	2-OH, 3-Me, 5-F
57	9	0	2-OH, 5-Cl
85	9	0	2-F
86	5	0	2-F
87	10	0	H
88	10	0	2-F
89	5	0	H
90	3	0	2-OCH ₃
91	3	0	2-CH ₃
92	3	0	2-F
93	3	0	H
94	9	0	2-OCH ₃
95	11	0	2-CH ₃
96	11	0	2-OCH ₃
97	11	0	2-F
98	11	0	H
99	9	0	2-CH ₃
100	9	0	H

101	5	0	2-CH ₃
102	1	0	2-OH, 4-OMe
103	2	0	2-OH
105	3	0	2-OH, 5-Cl
106	3	0	2-OH, 4-OMe
107	5	0	2-OH, 4-OMe
108	9	0	2-OH, 4-OMe
109	11	0	2-OH, 4-OMe
110	1	0	H
111	1	0	2-CH ₃
112	1	0	2-OMe
113	1	0	2-F
114	1	0	2-OH, 5-Cl
116	4	0	2-OH, 5-Cl
117	5	0	2-OH, 5-Cl
118	6	0	2-OH, 5-Cl

Cpd. 58:




Cpd. 59:




Cpd. 60: X=F


Cpd. 61: X=OH



Cpd. 62: X=SO

Cpd. 63: X=SO₂

Cpd. 64




Cpd. 65: X = Cl; Y = OMe

Cpd. 66: X = F; Y = OH


Cpd. 67: X = F; Y = OMe

Cpd. 68: X = OMe; Y = OH




Cpd. 69: n = 1


Cpd. 70: n = 2




Cpd. 71:




Cpd. 72:




Cpd. 73:




Cpd. 74: X=Me
 Cpd. 75: X=OMe




Cpd. 76: X = Y = OMe; Z = H; n = 1
 Cpd. 77: X = OH; Y = Z = Cl; n = 1




Cpd. 78 n=2
 Cpd. 134: n=3




Cpd. 79:




Cpd. 80:




Cpd. 81:



Cpd. 82:




Cpd. 83



Cpd. 84: $R = C(O)-NH-(CH_2)_7COOH$

Cpd. 135: $R = (CH_2)_7COOH$



The compositions of the present invention comprise at least one active agent, preferably a biologically or chemically active agent, and at least one of the compounds, or salts thereof, of the present invention. Methods for the preparation and administration of such compositions are also provided. The compositions comprising the compounds and active agents have utility in the delivery of active agents to selected biological systems and in an increased or improved bioavailability of the active agent compared to administration of the active agent alone.

DETAILED DESCRIPTION OF THE INVENTION

The compositions of the present invention include an active agent and a delivery agent. These compositions may be used to deliver various active agents through various biological, chemical, and physical barriers and are particularly suited for delivering active agents which are subject to environmental degradation.

Other advantages of the present invention include the use of easy to prepare, inexpensive raw materials. The compositions and the formulation methods of the present invention are cost effective, simple to perform, and amenable to industrial scale up for commercial production.

Compounds

The compounds may be in the form of the carboxylic acid and/or their salts. Salts include but are not limited to organic or inorganic salts, such as sodium salts. In addition, poly amino acids and peptides comprising one or more of these compound may be used.

An amino acid is any carboxylic acid having at least one free amine group and includes naturally occurring and synthetic

amino acids. Poly amino acids are either peptides (which are two or more amino acids joined by a peptide bond) or are two or more amino acids linked by a bond formed by other groups which can be linked by, e.g., an ester or an anhydride linkage. Peptides can vary in length from dipeptides with two amino acids to polypeptides with several hundred amino acids. See Chambers Biological Dictionary, editor Peter M.B. Walker, Cambridge, England: Chambers Cambridge, 1989, page 215. One or more of the amino acids or peptide units may be acylated or sulfonated.

Many of the compounds described herein may be derived from amino acids and can be readily prepared from amino acids by methods within the skill of those in the art based upon the present disclosure and the methods described in WO96/30036, WO97/36480, US 5,643,957 and US 5,650,386. For example, the compounds may be prepared by reacting the single amino acid with the appropriate acylating or amine-modifying agent, which reacts with a free amino moiety present in the amino acid to form amides. Protecting groups may be used to avoid unwanted side reactions as would be known to those skilled in the art. With regard to protecting groups, reference is made to T.W. Greene, Protecting Groups in Organic Synthesis, Wiley, New York (1981), the disclosure of which is hereby incorporated herein by reference.

The compound may be purified by recrystallization or by fractionation on one or more solid chromatographic supports, alone or linked in tandem. Suitable recrystallization solvent systems include, but are not limited to, acetonitrile, methanol, and tetrahydrofuran. Fractionation may be performed on a suitable chromatographic support such as alumina, using methanol/n-propanol mixtures as the mobile phase; reverse phase chromatography using trifluoroacetic acid/acetonitrile mixtures as the mobile phase; and ion exchange chromatography using water or an appropriate buffer as the mobile phase. When anion

exchange chromatography is performed, preferably a 0-500 mM sodium chloride gradient is employed.

Active Agents

Active agents suitable for use in the present invention include biologically active agents and chemically active agents, including, but not limited to, pesticides, pharmacological agents, and therapeutic agents.

For example, biologically or chemically active agents suitable for use in the present invention include, but are not limited to, proteins; polypeptides; peptides; hormones, and particularly hormones which by themselves do not pass (or which pass only a fraction of the administered dose) through the gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastro-intestinal tract; polysaccharides, and particularly mixtures of muco-polysaccharides; carbohydrates; lipids; other organic compounds; or any combination thereof.

Further examples include, but are not limited to, the following, including synthetic, natural or recombinant sources thereof: growth hormones, including human growth hormones (hGH), recombinant human growth hormones (rhGH), bovine growth hormones, and porcine growth hormones; growth hormone-releasing hormones; interferons, including α , β and γ ; interleukin-1; interleukin-2; insulin, including porcine, bovine, human, and human recombinant, optionally having counter ions including sodium, zinc, calcium and ammonium; insulin-like growth factor, including IGF-1; heparin, including unfractionated heparin, heparinoids, dermatans, chondroitins, low molecular weight heparin, very low molecular weight heparin and ultra low molecular weight heparin; calcitonin, including salmon, eel and human; erythropoietin; atrial natriuretic factor; antigens; monoclonal antibodies;

somatostatin; protease inhibitors; adrenocorticotropin, gonadotropin releasing hormone; oxytocin; leutinizing-hormone-releasing-hormone; follicle stimulating hormone; glucocerebrosidase; thrombopoietin; filgrastim; prostaglandins; cyclosporin; vasopressin; cromolyn sodium (sodium or disodium chromoglycate); vancomycin; desferrioxamine (DFO); parathyroid hormone (PTH), including its fragments; antimicrobials, including anti-fungal agents; analogs, fragments, mimetics or polyethylene glycol (PEG)-modified derivatives of these compounds; or any combination thereof.

Delivery systems

The compositions of the present invention comprise a delivery agent and one or more active agents. In one embodiment, one or more of the delivery agent compounds, or salts of these compounds, or poly amino acids or peptides of which these compounds or salts form one or more of the units thereof, may be used as a delivery agent by mixing with the active agent prior to administration.

The administration mixtures may be prepared by mixing an aqueous solution of the compound with an aqueous solution of the active ingredient, just prior to administration. Alternatively, the compound and the biologically or chemically active ingredient can be admixed during the manufacturing process.

The solutions may optionally contain additives such as phosphate buffer salts, citric acid, glycols, or other dispersing agents. Stabilizing additives may be incorporated into the solution, preferably at a concentration ranging between about 0.1 and 20% (w/v).

The delivery compositions of the present invention may also include one or more enzyme inhibitors. Such enzyme inhibitors include, but are not limited to, compounds such as actinonin or epiactinonin and derivatives thereof. Derivatives of these

compounds are disclosed in US 5,206,384. Other enzyme inhibitors include, but are not limited to, aprotinin (Trasylol) and Bowman-Birk inhibitor.

The amount of active agent is an amount effective to accomplish the purpose of the particular active agent for the target indication. The amount of active agent in the compositions typically is a pharmacologically, biologically, therapeutically, or chemically effective amount. However, the amount can be less than that amount when the composition is used in a dosage unit form, such as a capsule, a tablet, a powder, or a liquid, because the dosage unit form may contain a plurality of compound/biologically or chemically active agent compositions or may contain a divided pharmacologically, biologically, therapeutically, or chemically effective amount. The total effective amount can then be administered in cumulative units containing, in total, pharmacologically, biologically, therapeutically or chemically active amounts of biologically or pharmacologically active agent.

The total amount of active agent to be used can be determined by methods known to those skilled in the art. However, because the compositions may deliver active agents more efficiently than prior compositions, lower amounts of biologically or chemically active agents than those used in prior dosage unit forms or delivery systems can be administered to the subject, while still achieving the same blood levels and/or therapeutic effects.

The presently disclosed compounds deliver biologically and chemically active agents, particularly in oral, intranasal, sublingual, intraduodenal, subcutaneous, buccal, intracolonic, rectal, vaginal, mucosal, pulmonary, transdermal, intradermal, parenteral, intravenous, intramuscular and ocular systems, as well as traversing the blood-brain barrier.

Dosage unit forms can also include any one or combination of

excipients, diluents, disintegrants, lubricants, plasticizers, colorants, flavorants, taste-masking agents, sugars, sweeteners, salts, and dosing vehicles, including, but not limited to, water, 1,2-propane diol, ethanol, olive oil, or any combination thereof.

The compounds and compositions of the subject invention are useful for administering biologically or chemically active agents to any animals, including but not limited to birds such as chickens; mammals, such as cows, pigs, dogs, cats, primates, and particularly humans; and insects. The system is particularly advantageous for delivering chemically or biologically active agents which would otherwise be destroyed or rendered less effective by conditions encountered before the active agent reaches its target zone (i.e. the area in which the active agent of the delivery composition is to be released) and within the body of the animal to which they are administered. The compounds and compositions of the present invention are also useful in administering active agents, especially those which are not ordinarily deliverable by a particular route, especially by the oral route, or those for which improved delivery is desired. Delivery can be improved by delivering more active agent over a period of time, or in a particular time period (such as to effect quicker delivery).

Following administration, the active agent present in the composition or dosage unit form is taken up into the circulation.

The bioavailability of the agent is readily assessed by measuring a known pharmacological activity in blood, e.g. an increase in blood clotting time caused by heparin, or a decrease in circulating calcium levels caused by calcitonin. Alternately, the circulating levels of the active agent itself can be measured directly.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following examples illustrate the invention without limitation. All parts are given by weight unless otherwise indicated.

Example 1 - Compound Preparation

Method A: Preparation of compound 26. A 1L round bottom flask fitted with a magnetic stirrer was charged with 2-amino-p-cresol (1.71 g, 13.88 mmol, 1 equiv.) and 2M aqueous sodium hydroxide (20 ml). Methyl azeloyl chloride (3.08 g, 13.96 mmol, 1.01 eq.) Was added dropwise to the stirred solution at 0 C. After the addition, the reaction mixture was allowed to warm to ambient temperature and was stirred for 4-5 hours at ambient temperature. The pH of the solution was kept at about 11-12 by the addition of 20% sodium hydroxide. The solution was then extracted with ethyl acetate (3 x 30 ml). Combined organic layers were dried over sodium sulfate, filtered and concentrated in vacuo. The residue was redissolved in THF (50 ml) and treated with 2N NaOH (20 ml). The resulting mixture was stirred at room temperature for 8 hours. TLC indicated the reaction was complete. The mixture was concentrated in vacuo and acidified. The resulting solid was collected, recrystallized with methanol/acetone/water to afford 2.5 g of product.

Compounds 11-15, 18-21, 24, 25, 27-35, 48-50, 54-56, 58, 65-68, 73, 83, 110, 120 and 131-133 were also prepared by this method using the appropriate starting materials. Compounds 120, 122 and 123 were also prepared by this method using the appropriate starting materials, except that the reaction was run containing the THF.

Method B: Preparation of compound 60. A slurry of 1.03 g (5.62 mmol) of 3-amino-4-fluorohydrocinnamic acid and 20 ml of

methylene chloride was treated with 1.45 ml (1.24 g, 11.4 mmol) of trimethylsilyl chloride and was heated to reflux for 150 min.

The reaction mixture was cooled to 0C and treated with 2.4 ml (1.74 g, 17.2 mmol) of triethylamine. After stirring for 5 min, this mixture was poured into an addition funnel and added slowly to a 0C solution of 1.21 g (6.09 mmol) of O-acetylsalicyloyl chloride and 10 ml of methylene chloride. The reaction mixture was warmed to 25C and stirred for 18 hr. Washing with 3% aqueous hydrochloric acid (2 X 20 ml), water (1 X 20ml) and brine (1 X 15 ml), drying over sodium sulfate, and concentration in vacuo gave a brownish yellow solid. The solid was recrystallized from 15 ml of 65% ethanol/water to give a brown solid. The solid was dissolved in saturated aqueous sodium bicarbonate solution. After stirring until the acetate was cleaved off (by HPLC), the solution was acidified, causing a precipitate to develop. Isolation by filtration afforded 0.58 g of (38% yield) of product, mp = 202-4C.

Compounds 59, 61, 74-81, 107, 109 and 111-113 were also prepared by this method using the appropriate starting materials. Compounds 37, 42, 44 and 134 could also be prepared by this method using the appropriate starting materials.

Method C: Preparation of compound 1. 1,4-benzodioxan-2-one (4.50 g, 30 mmol) was dissolved in acetonitrile (75 ml) in a 250 ml round bottom flask equipped with a magnetic stir bar, argon purge and cold-water condenser. Triethylamine (4.17 ml, 3.03 g, 30 mmol) and 4-(4-aminophenyl)butyric acid (5.37 g, 30 mmol) were added. The reaction mixture was heated to reflux for 2 hr, stirred overnight at 25C and concentrated in vacuo. The brown residue was taken up in methylene chloride, and washed with aqueous 1N hydrochloric acid solution (1 X 100 ml). The solid that formed was isolated by filtration, and rinsed with aqueous 1N hydrochloric acid solution and water. The resulting solid was

dried overnight in a vacuum oven, yielding the compound as a tan solid (9.19 g, 93% yield).

Compound 53 was made by this method starting from dihydrocoumarin and 8-aminocaprylic acid.

Method D: Preparation of compound 36. 8-(N-3,5-dinitrosalicyloyl)aminocaprylic acid was prepared using Method T starting from 3,5-dinitrosalicylic acid and 8-aminocaprylic acid.

A solution of 0.7 g (1.9 mmol) of 8-(N-3,5-dinitrosalicyloyl)aminocaprylic acid and 40 ml of ethyl acetate was treated with 70 mg of 10% Pd/C and was placed under a hydrogen atmosphere for 18 hr. The catalyst was removed by filtration. The filtration was concentrated in vacuo. The residue was recrystallized from methanol/acetone/water to afford 0.4 g of (62% yield) of product, mp = 156-7C.

Method E: Preparation of compound 2. 4-(4-(4-benzyloxyphenoxyacetyl)amino)phenyl)butyric acid was prepared by the reaction of 4-(4-aminophenyl)butyric acid with 4-benzyloxyphenoxyacetyl chloride using Method C.

A slurry of 5.00 g (11.9 mmol) of 4-(4-(4-benzyloxyphenoxyacetyl)amino)phenyl)butyric acid and 150 ml methyl alcohol was stirred under argon for approximately 20 minutes. A catalytic amount (0.4 g) of 10% palladium on carbon was added in one portion. The reaction vessel was evacuated. The reaction was kept under a hydrogen atmosphere overnight at room temperature. The palladium on carbon was then filtered off and the filtrate was concentrated under vacuum to give the product as a white solid.

Compounds 8, 9 and 38 were also prepared by this method using the appropriate starting materials.

Method F: Preparation of compound 39. A suspension of 10.82 g (59.7 mmol) of 5-fluoroisatoic anhydride, 9.50 g of 8-aminocaprylic acid, 16.8 g of 50 wt % K_2CO_3 /water and 40 ml of dioxane was heated to reflux. After 2 hr, the reaction was complete (as measured by HPLC). The deep purple solution was cooled to 25C and acidified with 3 % aqueous hydrochloric acid to pH = 4.00, causing a dark precipitate to form. The solid was isolated by filtration and recrystallized from 65 % ethanol/water to give 11.86 g (67% yield) of the product as a yellow solid, mp = 108-9C.

Compounds 3-7, 10, 40 and 43 were also prepared by this method using the appropriate starting materials.

Method G: Preparation of compound 62. A 5C mixture of 4-(4-aminophenyl)butyric acid (1.0 eq) and aqueous 6N hydrochloric acid (5.44 eq) was treated with 1.05 eq of a 3N aqueous solution of sodium nitrite, adding slowly so as to keep the temperature below 5C. A solution of 2.8N aqueous potassium iodide (1.01 eq) was added. The reaction was stirred overnight. The layers were separated. The organic phase was purified by flash chromatography using methanol/methylene chloride as eluant to give 4-(4-iodophenyl)butyric acid.

A solution of 4-(4-iodophenyl)butyric acid (0.86M) and dimethylformamide was treated with 2 eq of potassium carbonate, 1.5 eq of 2-hydroxythiophenol and a catalytic amount of cuperic acetate (0.01 eq). The reaction mixture was refluxed for 18 hr, cooled to 25C, acidified with aqueous acid and extracted with ethyl acetate. The organic phase was concentrated. The residue was purified by flash chromatography using ethyl acetate/hexane as eluant to give 4-(4-(2-hydroxyphenyl)thiophenyl)butyric acid.

A 0.33M solution of 4-(4-(2-hydroxyphenyl)thiophenyl)butyric acid and ethyl acetate/acetic acid was treated with a 9.8M aqueous solution of hydrogen peroxide. After stirring for 12 hr,

the layers were separated. The organic phase was concentrated. Purification of the residue by flash chromatography using toluene/acetone/hexane as eluant gave the product.

Method H: Preparation of compound 82. A 0C solution of 3.97 g (17.8 mmol) of 9-bromo-1-nonanol and methylene chloride was treated with a solution of 2.91 g of 2-nitrophenylisocyanate and 10 ml of methylene chloride. The reaction mixture was heated to reflux for 2 hr, stirred at 25C for 16 hr and concentrated in vacuo. The yellow solid was identified as 9-bromononyl N-(2-nitrophenyl)carbamate and was used as is.

A suspension of 2.99 g (7.72 mmol) of 9-bromononyl N-(2-nitrophenyl)carbamate, 1.61 g (23.3 mmol) of sodium nitrite, 4.50 ml (4.72 g, 78.5 mmol) of acetic acid and 15 mL of dimethylsulfoxide was stirred at 35C for 7 hr. The reaction mixture was acidified with 3 % aqueous hydrochloric acid and extracted with diethyl ether (3 X 20 ml). The combined organic layers were extracted with 2N aqueous sodium hydroxide solution (3 X 20 ml). The basic aqueous phases were acidified with 3 % aqueous hydrochloric acid, causing a precipitate to develop. The solid was collected by filtration to give 0.79 g (30% yield) of compound, mp = 90-1C.

Method I: Preparation of compound 64. 4-(4-(2-aminobenzoyl)aminophenyl)butyric acid was prepared using Method F starting from isatoic anhydride and 4-(4aminophenyl)butyric acid.

A slurry of 4.73 g (16.0 mmole) 4-(4-(2-aminobenzoyl)aminophenyl)butyric acid and 40 ml of triethylorthoformate was placed under an argon atmosphere and heated to reflux for 18 hours. The reaction became clearer during the reflux. The reaction mixture was cooled to room

temperature and the resulting solid was collected by filtration to give 4.47g (88% yield) of the product, mp= 201-204C.

Method J: Preparation of compound 63. A 0.05M solution of 4-(4-(2-hydroxyphenyl)thiophenyl)butyric acid in methylene chloride was treated with 4 eq of m-chloroperbenzoic acid at 0C. The reaction mixture was allowed to warm to 25C and stirred for 12 hr. The solvent was stripped off. The residue was purified by flash chromatography using ethyl acetate/hexane/acetic acid as eluant to give the product.

Method K: Preparation of compound 84. 8-N-(2-aminobenzoyl)aminocaprylic acid was prepared using Method F, starting from isatoic anhydride and 8-aminocaprylic acid.

A slurry of 6.88 g (24.7 mmol) of 8-N-(2-aminobenzoyl)aminocaprylic acid and 100 mL of methylene chloride was cooled to 0C and treated with a solution of 2.00 ml (2.08 g, 12.4 mmol) of hexamethylenediisocyanate and 5 ml of methylene chloride. The reaction mixture was heated to reflux for 2 hr, cooled to 25C and diluted with 20 ml of ethanol. The resulting solid was isolated by filtration and recrystallized from 1/2/1 ethyl acetate/ethanol/water, yielding a total of 5.15 g (57% yield) of the product as a tan solid, mp = 138-142C.

Compound 135 could also be prepared by this method using the appropriate starting materials.

Method L: Preparation of compound 51. 8-(N-6-chloro-2-methoxybenzoyl)aminocaprylic acid was prepared using Method A, starting from 2-chloro-6-methoxybenzoic acid and 8-aminocaprylic acid.

A suspension of 1.27 g (3.72 mmol) of 8-(N-6-chloro-2-methoxybenzoyl)aminocaprylic acid in 200 ml of methylene chloride

under an argon atmosphere was cooled to 0C and treated with 8 ml of a 1.0 M boron tribromide solution in methylene chloride. After stirring for 60 min, TLC indicated that the reaction was complete. The reaction mixture was quenched with water and stirred for 30 min. The layers were separated. The aqueous layer was extracted with methylene chloride (2 X 30 ml). The combined organic layers were dried over sodium sulfate, filtered and concentrated in vacuo. The resulting white solid was recrystallized from methanol/acetone/water to afford 0.5 g (43% yield) of product, mp = 156-7C.

Method M: Preparation of compound 17. A solution of 4.28 mL (0.034 mole) of dihydrocoumarin, 75 ml of acetonitrile, 4.79 ml of triethylamine (3.48 g, 0.0343 mole), and 5.62 g of 3-(4-aminophenyl)propionic acid (0.034mole) was stirred under argon and the flask heated to reflux for 18 hours. The reaction was cooled to room temperature, and the acetonitrile removed under vacuum. The residue was stirred in methylene chloride and 1N aqueous hydrochloric acid, forming a white solid. The solid was filtered off, rinsed with water and methylene chloride, then vacuum dried at 50C to give 9.17g (86.1% yield) of product, mp=163-165C.

Compound 16 was also prepared by this procedure using the appropriate starting materials.

Method N: Preparation of compound 52. A solution of 10.0 g (65.8 mmol) of 3-hydroxyphenylacetic acid in 50 ml xylenes was treated with 6.45 ml (68.4 mmol) acetic anhydride. This mixture was refluxed for about 2.5 hours until most of the xylenes and acetic acid by-product was distilled off. The oligo-(3-hydroxyphenylacetic acid) was isolated as a brown oil.

This oil was dissolved in 150 ml of 1,4-dioxane. A solution

of 9.97 g (62.7 mmol) of 8-aminocaprylic acid and 34.5 ml of 2N NaOH solution was added to the oligomer solution. The reaction mixture was heated to reflux overnight. The dioxane was then removed under vacuum. The brown residue was taken up in 2N NaOH and extracted with two 100 ml portions of ethyl acetate. The aqueous layer was then acidified with 2N sulfuric acid solution and was then extracted with three 100 ml portions of ethyl acetate. The combined ethyl acetate layers were decolorized with activated carbon, dried with sodium sulfate, and concentrated under vacuum. The resulting brown oil was then purified by column chromatography using a silica gel column with ethyl acetate:hexane:acetic acid (60:40:1) as the mobile phase. The resulting white solid was washed with warm water (40-50C) to give the product as a white solid.

Compounds 41, 45 and 47 were also prepared by this method using the appropriate starting materials.

Method O: Preparation of compound 22. A solution of 3.0g (11.0 mmol) of 3(-4-(2 aminobenzoyl)aminophenyl)proprionic and 10 ml of methylene chloride. Acetic anhydride (1.12g, 1.04 ml, 11.0 mmol) was added dropwise over 30 minutes. Once the acetic anhydride was all added, the reaction was stirred at room temperature for 18 hr. The reaction was complete as determined by HPLC. The resulting solid was isolated by filtration. The resulting white solid was dried in a vacuum oven overnight yielding the product.

Method P: Preparation of compound 23. A mixture of 5.13 g (27.9 mmol) of 2-sulfobenzoic cyclic anhydride, 5.0 g (27.9 mmol) of 4-(4-aminophenyl)butyric acid and 100 ml of acetonitrile was stirred for 18 hr. The milky solution was concentrated. The residue was taken up into 50 ml of cold aqueous hydrochloric acid, extracted with ethyl acetate (5 X 50 ml) and concentrated.

The residue was purified by column chromatography using acetonitrile as the eluant to give the product.

Method Q: Preparation of compound 57. A solution of 10 g of 5-chloro-2-hydroxybenzamide (58.0 mmol), pyridine (22ml) and acetonitrile (25ml) was stirred in an ice bath. Ethyl chloroformate (6.1ml, 0.0638mol) was added dropwise. The pink solution was stirred for 30 minutes at <10°C. The ice bath was replaced with an oil bath. The reaction mixture was heated to 95°C and the volatiles were distilled off (43ml). The reaction was cooled to room temperature, causing a white solid to form. The mixture was poured into water (100ml) and acidified with concentrated aqueous hydrochloric acid. The resulting solid was filtered and recrystallized in hot ethanol to give 9.77g of 6-chloro-2H-1,3-benzoxazine-2,4(3H)-dione.

A slurry of 9.77 g of 6-chloro-2H-1,3-benzoxazine-2,4(3H)-dione (50 mmol) and 60 ml of DMF was treated with a solution of 1,10-dibromodecane (52.52g, 0.175mol) and DMF (60ml) was added. Diisopropylethylamine (9.6ml, 55 mmol) was added dropwise. A thermometer and condenser were attached, and the flask was placed in an oil bath. The reaction was heated to 60°C for approximately 3 hrs, cooled to 47°C and hexanes (150ml) were added. The mixture was diluted with water (175ml). The resulting solids were removed by filtration. The aqueous layer was washed with warm hexanes. The resulting solids were filtered from the hexane layer and recrystallized from warm hexanes to give 10.39 g of 6-chloro-3-(10-bromodecyl)-2H-1,3-benzoxazine-2,4(3H)-dione.

A mixture of 6-chloro-3-(10-bromodecyl)-2H-1,3-benzoxazine-2,4(3H)-dione (10.39g, 0.025mol), sodium nitrite (5.52g, 0.08mol) and DMSO (60ml) were stirred under argon. Acetic acid (14.9ml) was added and the reaction was heated to 75°C for 6 hrs. After being cooled to room temperature, the reaction mixture was then

dissolved in ethyl acetate and washed with 0.5N HCl (2x) and 2N NaOH (2x). The sodium hydroxide layers were stirred for 2 hrs and 20 minutes, and the solution was acidified with 2M H₂SO₄. The solids were then filtered and recrystallized from methylene chloride and hexanes. Yield: 3.0g.

Method R. Preparation of compound 71. A solution of 3.22 g (18.4 mmol) of mono-methyl phthlate, 2.90 ml (2.11 g, 20.8 mmol) of triethylamine and 20 ml of acetone was cooled in an ice (salt) bath and treated with a solution of 2.00 ml (2.27 g, 20.9 mmol) of ethyl chloroformate and 10 ml of acetone, added dropwise over 20 min. The white cloudy solution was stirred for 15 min and treated with a solution of 2.53 g (38.9 mmol) of sodium azide and 8 ml of water. After stirring for 30 min the still cloudy solution was poured into 50 ml of ice water and extracted with toluene (3 x 30 ml). The toluene phase was dried over sodium sulfate and heated to reflux for 80 min. The reaction mixture was cooled to 0C. In several portions, 3.11 g (18.8 mmol) of 3-(4-aminophenyl)propionic acid was added over 5 min. After stirring for 64 hr, the reaction mixture was diluted with 80 ml of a 3:1 dioxane/water solution and heated to reflux for 3 hr. The cooled mixture was extracted with 2N (1 x 30 ml) and 0.5N (2 x 30ml) sodium hydroxide. The combined aqueous layers were acidified, causing a white solid to form. Isolation by filtration, dissolution in ethyl acetate, filtration of the insoluble material and concentration of the filtrate gave 1.13 g of solid product.

Method S. Preparation of compound 69. A slurry of 3.17 g (21.7 mmol) of 4-hydroxyquinoazoline, 3 drops of dimethylformamide and 20.0 ml of thionyl chloride was heated to reflux. After 150 min, the clear yellow solution was cooled to 30C. The excess

thionyl chloride was distilled off at 50 mm vacuum. After 15 min at 1.0 mm vacuum, the off-white/yellow contents of the flask were taken up in 30 ml of methylene chloride and treated with a slurry of 3.61 g (21.9 mmol) of 3-(4-aminophenyl)propionic acid and 40 ml of isopropanol. The reaction mixture was stirred for 18 hr. The light yellow solid was isolated by filtration and dried in a vacuum oven to give the product.

Compound 70 was also prepared by this method except that acidification was done only to pH 4.56 so as to isolate the free amine.

Method T. Preparation of compound 46. Acetic anhydride (6.45 ml, 6.98 g, 68.4 mmol), 4-hydroxyphenylacetic acid (10.00 g, 65.8 mmol), and xylenes (50 ml) were added to a 100 mL, three-neck flask fitted with a magnetic stir bar, a thermometer, and a Dean-Stark trap with condenser. The flask was heated to reflux, the reaction mixture clearing to a yellow solution around 100°C. Most of the volatile organics (xylenes and acetic acid) were distilled into the Dean-Stark trap over two hours (135-146°C). Distillation was continued for another hour, during which the pot temperature slowly rose to 190°C and the distillate slowed to a trickle. The residue was poured off while still hot into an aluminum tray. Upon cooling a brown waxy solid formed.

A 2N sodium hydroxide (34.4 ml, 36 g, 68.7 mmol) and 8-aminocaprylic acid (9.94 g, 62.5 mmol) solution was added to a solution of oligo(4-hydroxyphenylacetic acid) (11.06 g, 81.3 mmol) and dioxane (150 ml), added over five minutes. The
5 reaction mixture was heated to 90°C for 5.5 hours (at which time the reaction was determined to have finished, by HPLC). The clear orange reaction mixture was cooled to 40°C. The dioxane was stripped off in vacuo. The brown residue was taken up in 2N sodium hydroxide, extracted with ethyl acetate (2 x 100 ml), and
10 acidified. Extraction with ethyl acetate (3 x 100 ml),

decolorizing with carbon, drying over sodium sulfate, and concentration in vacuo gave a brown oil. Trituration with warm water (2 times) and gave a tan solid which was recrystallized twice from ethanol/water to give 2.48 g of the product as a tan
5 solid.

Compounds 106 and 108 were also prepared by this method using the appropriate starting materials.

Method U. Preparation of compound 72. A suspension of 4.11
10 g (25.3 mmol) of 4-hydroxycoumarin, 4.54 g (25.3 mmol) of 4-(4-aminophenyl)butyric acid, and 20 ml of acetic acid was heated to reflux for 7 days. The reaction mixture was cooled to 25C, causing an off-white solid to form, which was collected by filtration. The filtrate was diluted with 50 ml of water,
15 causing a second solid to form, which was also collected by filtration. The two solids were combined and recrystallized from 65% ethanol/water to give 0.62 g of product.

Method V. Preparation of compound 85. A slurry of 5.00 g
20 of 10-aminodecanoic acid (26.7 mmol) in 70 mls methylene chloride was treated with 6.78 mls of chlorotrimethylsilane (5.80 g, 53.5 mmol) and was allowed to reflux for 140 min. The reaction mixture was cooled to 0C and was then treated with 5.58 mls triethylamine (4.1 g, 40.1 mmol). After this mixture stirred for
25 about 20 min, a solution of 3.91 mls of o-fluorobenzoyl chloride (4.24 g, 26.7 mmol) in 10 mls of methylene chloride was added dropwise to the reaction mixture over a period of 15 min. The reaction mixture was allowed to stir for 30 min. at 0C and then for 18 hrs at 25C. Methylene chloride was removed in vacuo and
30 100 mls of NaOH solution (2N) was added to the residue. This mixture was allowed to stir for 1 hr before the mixture was

acidified to pH=1 with hydrochloric acid solution (2M). The acidified mixture was then extracted with ethyl acetate (2 X 100 mls), decolorized with activated carbon, dried over sodium sulfate, and concentrated in vacuo. The resulting white solid
5 was recrystallized from a 50% ethanol/water mixture yielding a white solid, which was allowed to dry for 24 hr in vacuo at 25C. Yield of product was 6.57 g (79.5%), mp=85-86C.

Compounds 86-101 were also prepared by this method by reacting the appropriate amino acid with the appropriate acid
10 chloride.

Method W. Preparation of compound 102. A slurry of 20.72 g of glycine(0.276 mol) in 150 mls methylene chloride was treated with 70.06 mls of chlorotrimethylsilane (59.97 g, 0.552 mol) and
15 was allowed to reflux for 2 hours. The reaction mixture was cooled to 0C and was then treated with 115.41 mls triethylamine (83.79 g, 0.828 mol). After this mixture stirred for about 20 min, a solution of 20.72 g (0.276 mol) of 4-methoxy-2-acetylbenzoyl chloride (58.70 g, 0.276 mol) in 75 mls of
20 methylene chloride was added dropwise to the reaction mixture over a period of 15 min. The reaction mixture was allowed to stir for 30 min. at 0C and then for 18 hrs at 25C. Methylene chloride was removed in vacuo and 200 mls of NaOH solution (2N) was added to the residue. This mixture was allowed to stir for
25 several hours before the mixture was acidified to pH=3 with hydrochloric acid solution (2M). The resulting solids were filtered off and dried in vacuo at 40C. The solids were recrystallized from water/ethanol (3/1) yielding a solid, which was allowed to dry for 24 hr in vacuo at 25C. Yield of product
30 was 27.35 g (44%) mp= 185.5-189 C.

The sodium salt was made of the above solid by dissolving in 150 mls of ethanol with warming. Sodium hydroxide

(4.95 g in 14.5 mL of water) was added to the ethanol solution and cooled to room temperature. The resulting solid was filtered off using heptane to aid filtration and wash solids. After drying a tan solid was obtained (27.73, 92.37%) mp > 230 C. CHN
5 calc. for $C_{10}H_{10}N_1O_5Na \cdot 0.40H_2O$: C, 47.21; H, 4.28; N, 5.51; Na, 9.04; found: C, 47.14; H, 4.32; N, 5.36; Na, 8.45 and 2.83% water.

Method X. Preparation of compound 103. A slurry of 25.0 g
10 of β -alanine (0.281 mol) in 300 mls methylene chloride was treated with 71.33 mls of chlorotrimethylsilane (61.06 g, 0.562 mol) and was allowed to reflux for 1.5 hours. The reaction mixture was cooled to 0C and was then treated with 117.50 mls triethylamine (85.30 g, 0.843 mol). After this mixture stirred for about 20
15 min, a solution of acetylsalicyloyl chloride (55.73g, 0.281 mol.) in 150 mls of methylene chloride was added dropwise to the reaction mixture over a period of 15 min. The reaction mixture was allowed to stir for 30 min. at 0C and then for 18 hrs at 25C.
Methylene chloride was removed in vacuo and 200 mls of NaOH
20 solution (2N) was added to the residue. This mixture was allowed to stir for one hour before the mixture was acidified to pH=1 with sulfuric acid (2M). The resulting oil was extracted with ethyl acetate (3 x 200 mLs), dried over sodium sulfate and solvent removed in vacuo. The solids were recrystallized from
25 ethyl acetate/hexanes (1/1) yielding a solid, which was allowed to dry for 24 hr in vacuo at 25C. Yield of product was 9.20 g (16%). CHN calc. for $C_{10}H_{11}N_1O_4$: C, 57.03; H, 5.27; N, 6.67; found: C, 57.41; H, 5.30; N, 6.69. The sodium salt was made of the above solid by dissolving in 50 mls of ethanol with warming.
30 Sodium hydroxide (1.79 g in 5.25 mL of water) was added to the ethanol solution and cooled to room temperature. The resulting

solid was filtered off. After drying a solid was obtained (5.80 g) mp 231-235 C. CHN calc. for $C_{10}H_{10}N_1O_4Na \cdot 0.35H_2O$: C, 50.56; H, 4.54; N, 5.90; Na, 9.68; found: C, 50.30; H, 4.37; N, 5.72; Na, 9.55 and 2.68% water.

5

Method Y. Preparation of compound 104. 4-Methoxy salicyclic acid (98.74 g, 0.59 mol) was stirred in methylene chloride (500 ml) while in an ice bath. Triethylamine (123.4 ml, 1.5eq.) and acetyl chloride (46.2 ml, 1.1eq) were added dropwise respectively. The solution was removed from the ice bath and stirred at room temperature over two nights. The reaction was followed by HPLC. The reaction mixture was washed with 0.5 N HCl (2x200 ml) and water (2x200 ml). The organic layer was dried over sodium sulfate and reduce in vacuo. The resulting solid was recrystallized in methylene chloride/hexanes. The yield was 61.72 g of the 4-methoxy-2-acetyl benzoic acid. The structure was confirmed by 1H NMR.

To the mixture of 4-methoxy-2-acetyl benzoic acid (20g, 0.195 mol) and methylene chloride (100 ml), thionyl chloride (13.8 ml, 2 eq) and 1 drop DMF were added. The mixture refluxed for 1.5 hours, then cooled to room temperature and reduced in vacuo yielding an oil. The 4-methoxy-2-acetyl benzoyl chloride was used with no purification.

A slurry of 4-(4-aminophenyl)butyric acid (7.87 g, 0.044 mol) in methylene chloride (100 ml) was treated with TMS chloride (11.2 ml) and was allowed to reflux for 1.5 hours. The reaction mixture was cooled to 0° C and was then treated with triethylamine (18.4 ml) added dropwise. After this mixture was stirred for about 20 minutes, a solution of the 4-methoxy-2-acetylbenzoyl chloride (10 g) prepared above in methylene chloride (10 ml) was added dropwise to the reaction mixture over a period of 15 minutes. The ice bath was removed and the mixture

stirred at room temperature overnight. Methylene chloride was removed in vacuo and 2N NaOH solution (100 ml) was added to the residue. This mixture was allowed to stir for 1 hour before the mixture was acidified to pH=1 with 2M HCl. The acidified mixture
5 was then extracted with ethyl acetate (2x100 ml), decolorized with activated carbon, dried over sodium sulfate, and concentrated in vacuo. The resulting white solid was recrystallized from a 50% ethanol/water mixture yielding a white solid, which was allowed to dry for 24 hours in vacuo at 25C. CHN
10 calc. for $C_{18}H_{19}NO_5$: C, 65.54; H, 5.81; N, 4.25; found: C, 65.49; H, 5.84; N, 4.23. Yield was 7.76 g (53.6%), mp = 177-182C.

Method Z. Preparation of compound 105. 4-Bromobutyric acid (26.17 g, 0.16 mol) was added to methanol (150 ml) and several
15 drops sulfuric acid were added. This solution was refluxed for 3 1/4 hours. TLC (1:1 EtOAc/Hex) was done showing the completion of ester formation. The mixture reduced in vacuo to an oil. The oil was dissolved in methylene chloride and washed with water, saturated sodium bicarbonate, and brine. The organic layer was
20 dried over sodium sulfate and reduced in vacuo. The structure was confirmed by 1H NMR and yielded 20.26 g of methyl(4-bromo)butanoate.

6-Chlorocarsalam (12.4 g, 1.12eq), methyl(4-bromo)butanoate (10.13 g, 1.0 eq) and 10.13 g sodium carbonate (10.13 g, 1.12
25 eq) were stirred in 50 ml DMA. The solution was allowed to reflux for 4.5 hours, and then cooled to room temperature overnight. Solids were filtered off and washed with ethanol. Water and 2N NaOH were added to the filtrate. The mixture was heated for 2.5 hours. An HPLC was performed showing completion
30 of the hydrolysis. The solution was acidified with concentrated HCl to pH of approx. 1. The resulting white solid was filtered off, put over P_2O_5 in vacuo overnight. The solid was

recrystallized in methanol/water, filtered and dried yielding 7.35 g of compound. CHN calc. For $C_{11}H_{12}NO_4Cl$: C, 51.28; H, 4.69; N, 5.44; found: C, 50.92; H, 4.59; N, 5.46. Melting point 136-140C. Note that melting points of other batches of this compound made by the same method, showing purer product with elemental analysis, were in the range of 153-155C.

Method AA. Preparation of compound 118. Sodium carbonate (5.37g, 0.0506 mol) was added to 250 ml, 3-neck, round bottomed, flask containing 6-chlorocarsalam (10.0g, 0.0506 mol) and dimethylacetamide (50 ml). Ethyl- 5-bromoheptanoate (10.91g, 0.0460 mol) was added in one portion to the stirring reaction mixture, and heating of the reaction mixture was started. The reaction temperature was maintained at 80°C and allowed to heat for 16 hr. Heating was discontinued, and the reaction mixture was allowed to cool to room temperature. The reaction mixture was vacuum filtered and the filter cake was washed with two 20 ml portions of ethyl alcohol. Water was added to the filtrate until an orange-brown solid was noted to have precipitated. This solid was isolated by vacuum filtration and was washed first with 20 mls of ethyl alcohol and then with 20 mls of heptane. This solid was transferred to a round bottom flask and 200 ml of 2N NaOH was added. Heating to reflux was started and continued for one hr. The reaction was then cooled to 25 C and the reaction mixture was acidified with 2N HCl solution. The white solid, which precipitated, was isolated by filtration, was recrystallized from 30:70 ethyl alcohol:water and allowed to dry in vacuo overnight. 9.55g (63.0%) of the product was isolated. Melting point: 115-116C. Combustion analysis: %C: 56.09(calc.), 55.93(found); %H: 6.01(calc.), 6.09(found); %N:4.67(calc.), 4.64(found). 1H NMR Analysis: (d_6 -DMSO): δ 12.7, s, 1H (COOH); δ 12.0, s, 1H (OH); δ 8.88, t, 1H (NH); δ 7.94, d, 1H (H ortho to amide); δ

7.42, dd, 1H (H para to amide); δ 6.92, d, 1H (H ortho to hydroxide); δ 3.27, q, 2H (CH₂ alpha to amide); δ 2.20, t, 2H (CH₂ alpha to COOH); δ 1.40, m, 4H (CH₂ beta to amide, CH₂ beta to COOH); δ 1.30, m, 4H (remaining aliphatic CH₂).

- 5 Compounds 114, 116 and 117 were also prepared by this method using the appropriate starting materials.

Method BB: Preparation of compound 121. A suspension of 2 amino-4-chlorophenol (17.88g, 124.5 mmol), 8-ethoxy-8-oxooctanoic acid (25.19g, 124.5 mmol), boric acid (0.385g, 6.23 mmol), and 2-amino-5-methylpyridine (0.675g, 6.23 mmol) in 160 mL of dried toluene was heated at reflux (110C) under nitrogen for 4 hour during which water (2.5 mL) produced in the reaction was removed by azeotropic distillation in a Dean-Stark separation unit. Thin layer chromatography on silica gel (eluant: EtOAc/heptane: 1/1) revealed the complete reaction. To the cooled reaction mixture was added a 2N aqueous solution of NaOH (125 mL, 250 mmol). The reaction was heated at reflux for 4 h, then cooled. The cooled reaction mixture was diluted with ethyl acetate (300 mL) and water (150 mL). The aqueous layer was washed with two portions of ethyl acetate (250mL). After careful separation, the aqueous layer was chilled and acidified with a 10% solution of hydrochloric acid (86.20 mL, 250 mmol) to afford a solid which was filtered, washed with hexane, and dried under vacuum. Trituration with dichloromethane afforded the desired acid (8-(5-Chloro-2-hydroxyanilino)-8-oxooctanoic acid) (22.39g, 60%) as an off-white solid: HPLC (Column: Higgins Kromasil 100 C18, water/ acetonitrile/acetic acid: 950/50/1, 3mL/min, 220nm) R_t 5.38 min.; mp 123-124C; ¹H NMR (DMSO d₆ , 300 MHz) δ : 1.28 (m, 4H), 1.51 (m, 4H), 2.19 (t, 2H), 2.39 (t, 2H), 6.83 (d, 1H), 6.93 (dd, 1H), 7.95 (d, 1H), 9.20 (s, 1H), 10.10 (s, 1H), 12.00 (br

s, 1H); ^{13}C NMR (DMSO d_6 , 75 MHz) δ : 24.29, 24.90, 28.20, 33.58, 35.90, 116.50, 121.02, 122.20, 123.41, 127.74, 148.23, 171.93, 174.26. MS m/z 300 ($M+1$) $^+$. Anal. Calcd for $\text{C}_{14}\text{H}_{18}\text{ClNO}_4$: C, 56.10; H, 6.05; Cl, 11.83, N, 4.67. Found: C, 56.07, H, 6.11, 5 11.98, N, 4.64.

Compound 119 was prepared by the same method using the appropriate starting materials. Compounds 124-130 can also be prepared using this method with the appropriate starting materials.

10

Example 2 - Parathyroid Hormone Delivery

Oral gavage (PO) or intracolonic (IC) dosing solutions of delivery agent compound and parathyroid hormone residues 1-34 (PTH) (residues 1-38 for the solution with compound 103) were 15 prepared. A solution of the compound was made either with the sodium salt of the compound or by converting the free acid to its sodium salt by making a solution of the compound, stirring, adding one equivalent of sodium hydroxide (1.0 N), and diluting 20 with water (for PO solutions) or 25% aqueous propylene glycol (for IC). The final dosing solutions were prepared by mixing the compound solution with a PTH stock solution (typically having a concentration of 5 mg PTH/ml) and diluting to the desired volume (usually 3.0 ml). The compound and PTH dose amounts are listed 25 in Table 1 below.

Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and were administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The rats were administered one of the dosing solutions at 1 ml/kg for PO 30 or 0.5 ml/kg for IC. Blood samples were collected serially from the tail artery for serum determination of PTH concentration. Serum PTH concentrations were quantified by a PTH

radioimmunoassay kit (Kit # RIK 6101 from Peninsula Laboratories, Inc., San Carlos, CA). Results are illustrated in Table 1, below.

Table 1. PTH Delivery in Rats - oral (PO) and intracolonic (IC)				
Compound	Method of Administration	Compound Dose (mg/kg)	PTH Dose (μ g/kg)	Mean Peak Serum PTH (pg/mL)
3	PO	300	100	222 \pm 155
10	PO	300	100	420 \pm 335
79	IC	100	25	731 \pm 577
80	IC	100	25	1456 \pm 486
86	PO	100	200	0
89	PO	100	200	27 \pm 61
90	PO	100	200	14 \pm 21
91	PO	100	200	5 \pm 12
92	PO	100	200	303 \pm 427
93	PO	100	200	343 \pm 155
94	PO	100	200	17 \pm 38
102	PO	100	200	252.13 \pm 230.46
102	PO	100	200	70.98 \pm 81.81
102	PO	100	200	894.82 \pm 702.01
102	PO	100	200	185.52 \pm 59.47
103	IC	100	25	38.53 \pm 30.9
104	PO	100	200	286.35 \pm 191.58
106	PO	100	200	309.07 \pm 289.74
106	PO	100	200	894.91 \pm 1220.06
106	PO	100	200	1459.71 \pm 1041.36
106	PO	100	200	192.15 \pm 48.81
107	PO	100	200	110.19 \pm 142.23
107	PO	100	200	254.71 \pm 191.97
107	PO	100	200	1302.99 \pm 871.82
107	PO	100	200	304.8 \pm 381.39

Example 3 - Heparin Delivery

Intracolonic dosing (IC) compositions containing delivery agent compound and heparin sodium USP in 25% aqueous propylene glycol were prepared by mixing. Either the sodium salt of the compound was used or the free acid was converted to the sodium salt with one equivalent of sodium hydroxide (1.0 N). Typically, compound and heparin powders were mixed, 25% aqueous propylene glycol was added, NaOH solution was added, the contents sonicated, then diluted to a volume of 3.0. pH was checked and adjusted if necessary to pH=7-8. The final compound and heparin dose amounts are listed in Table 2 below.

Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and were administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The dosing solutions were administered to fasted rats at a volume dose of 1 ml/kg. Blood samples were collected by cardiac puncture following the administration of ketamine (44 mg/kg). Heparin activity was determined utilizing the activated partial thromboplastin time (APTT) according to the method of Henry, J.B., Clinical Diagnosis and Management by Laboratory Methods; Philadelphia, PA; W.B. Saunders (1979). Results are given in Table 2, below.

Table 2. Intracolonic Delivery of Heparin

Compound	Method of Administration	Compound Dose (mg/kg)	Heparin Dose (mg/kg)	Mean Peak APTT (sec)
7	IC	50	25	59 ± 41
14	IC	50	25	54 ± 21
28	IC	50	25	55 ± 27
33	IC	50	25	42 ± 21
34	IC	50	25	58 ± 31

Table 2. Intracolonic Delivery of Heparin

Compound	Method of Administration	Compound Dose (mg/kg)	Heparin Dose (mg/kg)	Mean Peak APTT (sec)
35	IC	50	25	154 ± 171
41	IC	50	25	41 ± 26
46	IC	50	25	52 ± 34
48	IC	50	25	75 ± 18
51	IC	50	25	111 ± 49
54	IC	50	25	124 ± 137
55	IC	50	25	125 ± 195
56	IC	50	25	91 ± 75
60	IC	50	25	71 ± 43
72	IC	50	25	50 ± 18
85	IC	50	25	27 ± 4
86	IC	50	25	24 ± 1
86	IC	50	25	31 ± 11
87	IC	50	25	21 ± 1
87	IC	50	25	23 ± 3
88	IC	50	25	59 ± 47
89	IC	50	25	33 ± 7
90	IC	50	25	26 ± 7
91	IC	50	25	24 ± 4
92	IC	50	25	22 ± 2
93	IC	25	50	22 ± 0
94	IC	50	25	50 ± 28
95	IC	50	25	30 ± 2
96	IC	50	25	72 ± 63
97	IC	50	25	33 ± 10
98	IC	50	25	25 ± 5
99	IC	50	25	34 ± 7
100	IC	50	25	31 ± 8
101	IC	50	25	26 ± 5
102	IC	50	25	24.8 ± 0.9
102	IC	50	25	24.7 ± 6.5
103	IC	50	25	21.9 ± 2.0
106	IC	50	25	48 ± 16.9

Table 2. Intracolonic Delivery of Heparin

Compound	Method of Administration	Compound Dose (mg/kg)	Heparin Dose (mg/kg)	Mean Peak APTT (sec)
106	IC	50	25	27.7 ± 12.6
107	IC	50	25	26.2 ± 6.1
108	IC	50	25	72.9 ± 28.9
109	IC	50	25	24.2 ± 1.7
110	IC	50	25	26.5 ± 4.7
110	IC	50	25	23.4 ± 0.7
111	IC	50	25	24.4 ± 3.3
112	IC	50	25	28.7 ± 11.5
113	IC	50	25	20.4
120	IC	50	25	42 ± 34
131	IC	50	25	58 ± 30
132	IC	50	25	65 ± 19

Example 4: Recombinant Human Growth Hormone (rhGH) Oral Delivery

Oral gavage (PO) dosing solutions of delivery agent compound and rhGH in phosphate buffer were prepared. A solution of the compound was made either with the sodium salt of the compound or by converting the free acid to its sodium salt by making a solution of the compound, stirring, adding one equivalent of sodium hydroxide (1.0 N), and diluting with phosphate buffer. The final dosing solutions were prepared by mixing the compound solution with an rhGH stock solution (typically having a concentration of 15 mg rhGH/ml) and diluting to the desired volume (usually 3.0 ml). The compound and rhGH dose amounts are listed in Table 3 below.

Male Sprague-Dawley rats weighing 200-250 g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The rats were administered 1 ml/kg of the dosing solution. Blood samples were

collected serially from the tail artery for determination of serum rhGH concentrations. Serum rhGH concentrations were quantified by an rhGH immunoassay test kit (Kit # K1F4015 from Genzyme Corporation Inc., Cambridge, MA).

5 Results are presented in Table 3 below.

Table 3. Oral Delivery of rhGH in Rats				
Compound	Method of Administration	Compound Dose (mg/kg)	rhGH Dose (mg/kg)	Mean Peak Serum [rhGH] (ng/ml)
3	PO	300	6	72 ± 45
10	PO	200	3	43 ± 65
40	PO	300	6	42 ± 80
45	PO	200	3	49 ± 56
54	PO	200	3	48 ± 33
74	PO	200	3	80 ± 44
76	PO	200	3	40 ± 34
77	PO	200	3	54 ± 62

10 Example 5 Salmon Calcitonin (sCT) Delivery

Oral dosing (PO) compositions of delivery agent compound and salmon calcitonin (sCT) in water were prepared by mixing. The amounts are listed in Table 4. 450 mg of compound was added to 2.0 ml of water. Either the sodium salt of the compound was used or the free acid was converted to the sodium salt by stirring the resultant solution and adding one equivalent of sodium hydroxide (1.0 N) and diluting with water. 90 µg sCT was added to the solution. Water was then added to bring the total volume to 3.0 ml. The solution had a final compound concentration of 150 mg/ml. (For compounds 118 and 123, the solutions were diluted to 6.0 ml and the volume dose was doubled.) The total sCT

concentration was 30 µg/ml. (For compound 123, a different amount of sCT was used to result in the final sCT dose of 100 µg/kg when 2.0 ml/kg was dosed.)

Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and were administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The rats were administered 1 ml/kg of dosing solution (2 ml/kg for compounds 118 and 123). Blood samples were collected serially from the tail artery. Serum sCT was determined by testing with an EIA kit (Kit # EIAS-6003 from Peninsula Laboratories, Inc., San Carlos, CA). The standard protocol from the kit was modified for compounds 104 and 105 as follows: incubated with 50 µl peptide antibody for 2 hours with shaking in the dark, washed the plate, added serum and biotinylated peptide and diluted with 4 ml buffer, and shook overnight in the dark. Results are illustrated in Table 4, below.

Table 4. Oral Delivery of Salmon Calcitonin (sCT) in Rats			
<u>Compound</u>	<u>Compound dose (mg/kg)</u>	<u>sCT Dose (µg/kg)</u>	<u>Serum sCT (pg/ml) ± SD (SE)</u>
104	50	25	287 ± 104
105	50	25	583 ± 140
105	150	30	802 ± 669 (299)
116	150	30	724 ± 463 (207)
117	150	30	383 ± 292 (131)
118	150	30	276 ± 319 (159)
119	150	30	95 ± 119 (53)
121	150	30	717 ± 603 (301)
122	150	30	187 ± 79 (36)
123	150	100	0

The above mentioned patents, applications, test methods, and publications are hereby incorporated by reference in their entirety.

- 5 Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. All such obvious variations are within the fully intended scope of the appended claims

WHAT IS CLAIMED IS:

1. A compound selected from the group consisting of compounds 1-135 and salts thereof.

2. A composition comprising:

(A) an active agent; and

(B) a compound selected from the group consisting of compounds 1-135, salts thereof, and mixtures thereof.

5

3. The composition of claim 2, wherein the active agent is selected from the group consisting of a biologically active agent, a chemically active agent, and a combination thereof.

4. The composition of claim 3, wherein the biologically active agent comprises at least one protein, polypeptide, peptide, hormone, polysaccharide, mucopolysaccharide, carbohydrate, or lipid.

5

5. The composition of claim 3, wherein the biologically active agent is selected from the group consisting of:

growth hormones, human growth hormones (hGH), recombinant human growth hormones (rhGH), bovine growth hormones, porcine

5 growth hormones, growth hormone-releasing hormones, interferons,
 α -interferon, β -interferon, γ -interferon, interleukin-1,
interleukin-2, insulin, porcine insulin, bovine insulin, human
insulin, and human recombinant insulin, insulin-like growth
factor (IGF), IGF-1, heparin, unfractionated heparin,
10 heparinoids, dermatans, chondroitins, low molecular weight
heparin, very low molecular weight heparin, ultra low molecular
weight heparin, calcitonin, salmon calcitonin, eel calcitonin,
human calcitonin; erythropoietin (EPO), atrial natriuretic factor,
antigens, monoclonal antibodies, somatostatin, protease
15 inhibitors, adrenocorticotropin, gonadotropin releasing hormone,
oxytocin, leutinizing-hormone-releasing-hormone, follicle
stimulating hormone, glucocerebrosidase, thrombopoietin,
filgrastim, prostaglandins, cyclosporin, vasopressin, cromolyn
sodium, sodium chromoglycate, disodium chromoglycate, vancomycin,
20 desferrioxamine (DFO), parathyroid hormone (PTH), fragments of
PTH, antimicrobials, anti-fungal agents; analogs, fragments,
mimetics and polyethylene glycol (PEG)-modified derivatives of
these compounds; and any combination thereof.

6. The composition of claim 3, wherein the biologically
active agent comprises insulin, unfractionated heparin, low
molecular weight heparin, very low molecular weight heparin,
ultra low molecular weight heparin, calcitonin, parathyroid

5 hormone, erythropoietin, human growth hormones, or combinations thereof.

7. A dosage unit form comprising:

(A) the composition of claim 2; and

(B) (a) an excipient

(b) a diluent,

5 (c) a disintegrant,

(d) a lubricant,

(e) a plasticizer,

(f) a colorant,

(g) a dosing vehicle, or

10 (h) any combination thereof.

8. The dosage unit form of claim 7, wherein the active agent is selected from the group consisting of a biologically active agent, a chemically active agent, and a combination thereof.

5

9. The dosage unit form of claim 8, wherein the biologically active agent comprises at least one protein, polypeptide, peptide, hormone, polysaccharide, mucopolysaccharide, carbohydrate, or lipid.

5

10. The dosage unit form of claim 8, wherein the biologically active agent is selected from the group consisting of:

growth hormones, human growth hormones (hGH), recombinant
5 human growth hormones (rhGH), bovine growth hormones, porcine
growth hormones, growth hormone-releasing hormones, interferons,
 α -interferon, β -interferon, γ -interferon, interleukin-1,
interleukin-2, insulin, porcine insulin, bovine insulin, human
insulin, and human recombinant insulin, insulin-like growth
10 factor(IGF), IGF-1, heparin, unfractionated heparin, heparinoids,
dermatans, chondroitins, low molecular weight heparin, very low
molecular weight heparin, ultra low molecular weight heparin,
calcitonin, salmon calcitonin, eel calcitonin, human calcitonin;
erythropoietin (EPO), atrial natriuretic factor, antigens,
15 monoclonal antibodies, somatostatin, protease inhibitors,
adrenocorticotropin, gonadotropin releasing hormone, oxytocin,
luteinizing-hormone-releasing-hormone, follicle stimulating
hormone, glucocerebrosidase, thrombopoietin, filgrastim,
prostaglandins, cyclosporin, vasopressin, cromolyn sodium, sodium
20 chromoglycate, disodium chromoglycate, vancomycin,
desferrioxamine (DFO), parathyroid hormone (PTH), fragments of
PTH, antimicrobials, anti-fungal agents; analogs, fragments,
mimetics and polyethylene glycol (PEG)-modified derivatives of
these compounds; and any combination thereof.

25

11. The dosage unit form of claim 8, wherein the biologically active agent comprises insulin, unfractionated heparin, low molecular weight heparin, very low molecular weight heparin, ultra low molecular weight heparin, calcitonin, parathyroid hormone, erythropoietin, human growth hormones, or combinations thereof.

12. The dosage unit form of claim 7, wherein the dosage unit form comprises a dosing vehicle comprising a tablet, a capsule, a powder, or a liquid.

13. The dosage unit form of claim 7, wherein the dosing vehicle is liquid selected from the group consisting of water, 1,2-propane diol, ethanol, and any combination thereof.

14. A method for administering an active agent to an animal in need of the agent, the method comprising administering orally to the animal the composition of claim 2.

15. A method for preparing a composition comprising mixing:

- (A) at least one active agent;
- (B) the compound of claim 1; and

5

(C) optionally, a dosing vehicle.

16. A compound comprising compound 105 and salts thereof.

17. A composition comprising:

(A) an active agent; and

(C) the compound of claim 16.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 233/25, 237/36, 237/40, A61K 47/18	A3	(11) International Publication Number: WO 00/07979 (43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: PCT/US99/17974 (22) International Filing Date: 6 August 1999 (06.08.99) (30) Priority Data: 60/095,778 7 August 1998 (07.08.98) US 60/098,500 31 August 1998 (31.08.98) US 60/108,366 13 November 1998 (13.11.98) US 60/119,207 5 February 1999 (05.02.99) US (71) Applicant (for all designated States except US): EMISPHERE TECHNOLOGIES, INC. [US/US]; 765 Old Saw Mill River Road, Tarrytown, NY 10591 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GSCHNEIDNER, David [US/US]; 44 Cerretta Street #6, Stamford, CT 06907 (US). LEONE-BAY, Andrea [US/US]; 20 Woodland Way, Ridgefield, CT 06877 (US). WANG, Eric [US/US]; 8205 Featherhill Road #302, Perryhall, MD 21128 (US). ERRIGO, Lynn [US/US]; 85 Birch Street, Port Chester, NY 10573 (US). KRAFT, Kelly [US/US]; 38 1/2 Wolden Road, Ossining, NY 10562 (US). MOYE-SHERMAN, Destardi [US/US]; 102 Chateau Rive, Peekskill, NY 10566 (US). HO, Koc-Kan [MC/US]; 70 Barker Street, Mount Kisco, NY 10549 (US). PRESS, Jeffrey, Bruce [US/US]; Bear Berry Lane, Brew-		ster, NY 10509 (US). WANG, Nai, Fang [CN/US]; Apartment 2C, 41-42 42nd Street, Long Island City, NY 11104 (US). (74) Agents: LESSLER, Jay, P. et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022-7513 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 18 May 2000 (18.05.00)
(54) Title: COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS (57) Abstract Compounds and compositions for the delivery of active agents are provided. Methods of administration and preparation are provided as well.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/17974

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07C233/25 C07C237/36 C07C237/40 A61K47/18		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07C A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 12473 A (DEMORIN FRENEL ;HO KOK KAN (US); PATON DUNCAN R (US); EMISPHERE TE) 2 May 1996 (1996-05-02) the whole document	1-15
Y	WO 96 12474 A (EMISPHERE TECH INC ;SANTIAGO NOEMI B (US); HAAS SUSAN (US); LEONE) 2 May 1996 (1996-05-02) the whole document	1-15
Y	WO 96 12475 A (BARANTSEVITCH EVGUENI ;WANG NAI FANG (US); MILSTEIN SAM J (US); EM) 2 May 1996 (1996-05-02) the whole document	1-15
-/--		
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-size: 1.2em;">7 December 1999</div>		Date of mailing of the international search report <div style="text-align: center; font-size: 1.2em;">24. 03. 00</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-size: 1.2em;">Janus, S</div>

INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/US 99/17974

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 21464 A (EMISPHERE TECH INC ;SANTIAGO NOEMI B (US); HAAS SUSAN (US); LEONE) 18 July 1996 (1996-07-18) the whole document ---	1-15
Y	WO 96 30036 A (EMISPHERE TECH INC ;LEONE BAY ANDREA (US); HO KOC KAN (US); SARUBB) 3 October 1996 (1996-10-03) the whole document ---	1-15
Y	WO 97 36480 A (EMISPHERE TECHNOLOGIES INC ;LEONE BAY ANDREA (US)) 9 October 1997 (1997-10-09) the whole document ---	1-15
Y	WO 98 21951 A (HAAS SUSAN ;MILSTEIN SAM J (US); EMISPHERE TECH INC (US)) 28 May 1998 (1998-05-28) the whole document ---	1-15
P,Y	WO 98 34632 A (GSCHNEIDNER DAVID ;WANG ERIC (US); HO KOC KAN (US); LEIPOLD HARRY) 13 August 1998 (1998-08-13) the whole document ---	1-15
L,E	WO 98 49135 A (EMISPHERE TECH INC) 5 November 1998 (1998-11-05) the whole document ---	1-15
P,Y	WO 99 16427 A (LEIPOLD HARRY ;MILSTEIN SAM J (US); EMISPHERE TECH INC (US); LEONE) 8 April 1999 (1999-04-08) the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

II. International application No.

PCT/US 99/ 17974

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1 - 15 (in part)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/17974

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-15 (in part)

Compounds 1-25, 73-75, 79-81, 104 and 133, and their use as targeting agents.

2. Claims: 1-15 (in part)

Compounds 26-35, 83 and 119-132, and their use as targeting agents.

3. Claims: 1-15 (in part), 16 and 17

Compounds 36-53, 65-68, 85-103 and 105-118, and their use as targeting agents.

4. Claims: 1-15 (in part)

Compound 59 and its use as targeting agent.

5. Claims: 1-15 (in part)

Compound 60, 61, 76 and 77, and their use as targeting agents.

6. Claims: 1-15 (in part)

Compounds 62 and 63 and their use as targeting agent.

7. Claims: 1-15 (in part)

Compound 64 and its use as targeting agent.

8. Claims: 1-15 (in part)

Compounds 69 and 70 and their use as targeting agents.

9. Claims: 1-15 (in part)

Compound 71 and its use as targeting agent.

10. Claims: 1-15 (in part)

Compound 72 and its use as targeting agent.

11. Claims: 1-15 (in part)

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99/17974

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Compounds 78 and 134 and their use as targeting agents.

12. Claims: 1-15 (in part)

Compound 82 and its use as targeting agent.

13. Claims: 1-15 (in part)

Compounds 84 and 135 and their use as targeting agents.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 99/17974

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9612473 A	02-05-1996	US 5643957 A	01-07-1997
		AU 711887 B	21-10-1999
		AU 3963395 A	15-05-1996
		BR 9510168 A	14-10-1997
		CA 2203033 A	02-05-1996
		CZ 9701229 A	15-10-1997
		EP 0783299 A	16-07-1997
		FI 971776 A	25-04-1997
		HU 77759 A	28-07-1998
		JP 10507762 T	28-07-1998
		NO 971889 A	23-06-1997
		PL 319833 A	01-09-1997
		US 5955503 A	21-09-1999
WO 9612474 A	02-05-1996	US 5958457 A	28-09-1999
		AU 4005295 A	15-05-1996
		CA 2202299 A	02-05-1996
		EP 0784469 A	23-07-1997
		JP 10509432 T	14-09-1999
WO 9612475 A	02-05-1996	US 5714167 A	03-02-1998
		AU 4152496 A	15-05-1996
		EP 0781124 A	02-07-1997
		JP 10509433 T	14-09-1998
WO 9621464 A	18-07-1996	US 5709861 A	20-01-1998
		AU 704776 B	06-05-1999
		AU 5131596 A	31-07-1996
		CA 2209492 A	18-07-1996
		EP 0812212 A	17-12-1997
		FI 972945 A	11-07-1997
		HU 9702307 A	30-03-1998
		JP 11514331 T	07-12-1999
		NO 973250 A	05-09-1997
		PL 321275 A	24-11-1997
WO 9630036 A	03-10-1996	US 5650386 A	22-07-1997
		AU 712222 B	04-11-1999
		AU 5662996 A	16-10-1996
		BR 9604880 A	19-05-1998
		CA 2214323 A	03-10-1996
		CN 1190893 A	19-08-1998
		CZ 9703073 A	18-03-1998
		EP 0817643 A	14-01-1998
		FI 973828 A	29-09-1997
		HU 9901162 A	30-08-1999
		NO 974495 A	28-11-1997
		NZ 307319 A	28-10-1999
		PL 322494 A	02-02-1998
		US 5965121 A	12-10-1999
		US 5989539 A	23-11-1999
		US 5866536 A	02-02-1997
		US 6001347 A	14-12-1999
WO 9736480 A	09-10-1997	AU 2595697 A	22-10-1997
		AU 712222 B	04-11-1999
		EP 0817643 A	14-01-1998
		SI 9720025 A	31-08-1999

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 99/17974

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9736480 A		US 5650386 A	22-07-1997
		US 5965121 A	12-10-1999
		US 6001347 A	14-12-1999
		US 5989539 A	23-11-1999
		US 5866536 A	02-02-1999
WO 9821951 A	28-05-1998	CA 2243643 A	28-05-1998
		EP 0886471 A	30-12-1998
WO 9834632 A	13-08-1998	US 5776888 A	07-07-1998
		US 5804688 A	08-09-1998
		US 5876710 A	02-03-1999
		US 5939381 A	17-08-1999
		US 5990166 A	23-11-1999
		US 5773647 A	30-06-1998
		US 5879681 A	09-03-1999
		AU 6275698 A	26-08-1998
WO 9849135 A	05-11-1998	US 5863944 A	26-01-1999
		AU 6959098 A	24-11-1998
		EP 0979225 A	16-02-2000
WO 9916427 A	08-04-1999	AU 9513698 A	23-04-1999